

12

# EUROPEAN PATENT APPLICATION

21 Application number: 89313429.6

51 Int. Cl.<sup>5</sup>: **C07C 227/32, C07C 229/22, B01J 31/24**

22 Date of filing: 21.12.89

30 Priority: 22.12.88 JP 324651/88

43 Date of publication of application: 27.06.90 Bulletin 90/26

64 Designated Contracting States:  
CH DE FR GB IT LI NL


71 Applicant: Takasago International Corporation  
19-22, Takanawa 3-chome Minato-ku  
Tokyo(JP)

72 Inventor: Taketomi, Takanao  
112-7, Ookido-cho  
Chiba-shi Chiba(JP)  
Inventor: Sakaguchi, Toshiaki  
3-3-35, Minamihara  
Hiratsuka-shi Kanagawa(JP)  
Inventor: Kumobayashi, Hidenori  
1-4-39, Nakakaigan  
Chigasaki-shi Kanagawa(JP)


74 Representative: Moore, Anthony John et al  
Gee & Co. Chancery House Chancery Lane  
London WC2A 1QU(GB)

54 Process for preparing optically active carnitine ester.

57 A process for preparing an optically active carnitine ester represented by formula (I):



wherein R is lower alkyl and X is halogen, comprises asymmetrically hydrogenating a  $\gamma$ -trimethylammonium-3-oxobutanoic ester halide represented by formula (II):



wherein R and X are as defined above, in the presence of a ruthenium-optically active phosphine complex as a catalyst. An optically active carnitine ester of any desired isomerism can be obtained through simple operation in high yield at high optical purity. The substrate used is readily available.

EP 0 375 417 A2

## PROCESS FOR PREPARING OPTICALLY ACTIVE CARNITINE ESTER

This invention relates to a process for preparing an optically active carnitine ester which is an intermediate for obtaining medicinally important optically active carnitine. More particularly, it relates to a process for preparing an optically active carnitine ester by asymmetric hydrogenation of a  $\gamma$ -trimethylammonium-3-oxobutanoic ester halide in the presence of a ruthenium-optically active phosphine complex as a catalyst.

L-Carnitine is present in a living body and functions to carry activated long-chain free fatty acids through the mitochondrial inner membrane. Acyl-CoA derivatives are allowed to enter into the mitochondrial matrix through the mitochondrial inner membrane only in the form of its ester with L-carnitine. Such a carrier function of L-carnitine is exerted in delivering the activated long-chain fatty acids from the site of their biosynthesis. The carnitine found in the living body is exclusively levorotatory, i.e., of L-form, while D-carnitine has never been detected in living bodies.

On the other hand, racemic carnitine has been used for years for various purposes. For example, DL-carnitine has been sold in Europe as an appetizer and also reported to have an effect of promoting physical development of children as described in Borniche et al., *Olinioa Chemica Acta*, Vol. 5, pp. 171 to 176 (1960). However, in recent years, the importance of using only L-carnitine for therapeutic purposes has been increasing. That is, D-carnitine has turned out to antagonize against carnitine acyltransferase, e.g., carnitine acetyl transferase (CAT) and carnitine palmitoyl transferase (PTC). It has also been elucidated recently that D-carnitine causes depletion of L-carnitine in cardiac tissues. Accordingly, it is essential that cases under treatment for cardiac diseases or for reduction of blood fat should receive only L-carnitine. L-Carnitine has thus been recognized as an important medicine.


Known techniques for preparing an optically active carnitine ester and carnitine include (i) optical resolution of a racemate, (ii) biochemical reduction of the corresponding keto ester, and (iii) synthesis using other optically active substances as a starting material or an intermediate.

Examples of the technique (i) include a process in which DL-carnitine amide hydrochloride is subjected to an ion exchange treatment and then resolved using D-camphoric acid as disclosed in JP-A-55-13299 (the term "JP-A" as used herein means an "unexamined published Japanese patent application") and a process in which DL-carnitine nitrile is resolved using L-camphor-10-sulfonic acid as disclosed in JP-B-40-3891 (the term "JP-B" as used herein means an "examined published Japanese patent application"). Examples of the technique (ii) include a process in which a  $\gamma$ -haloacetoacetic ester is converted to an optically active  $\gamma$ -halo- $\beta$ -hydroxybutyric ester by microbial fermentation and then quaternarized with trimethylamine as disclosed in JP-A-59-118093 and a process in which  $\gamma$ -azidoacetoacetic ester is reduced with a microorganism to obtain an optically active  $\gamma$ -azido- $\beta$ -hydroxybutyric ester. Examples of the technique (iii) include a process using D-mannitol as disclosed in JP-A-57-165352 and a process using (S)-chloromethyloxirane as disclosed in JP-A-62-212382.


However, the processes (i) utilizing resolution of a racemate as a carnitine precursor require a resolving agent in an amount equimolar to the substrate. Also, the highest possible yield of the desired product, attained if the reaction proceeds ideally, is 50%. Also, the undesired enantiomer is useless or must be racemized for re-use. The processes (ii) utilizing microbial asymmetric reduction are disadvantageous in that the usable substrate is limited, the production efficiency is low in many cases, and isolation of the product from the reaction solution involves complicated procedures. The processes (iii) using a naturally-occurring optically active substance as a starting material have disadvantages, such as requirement of long reaction steps.

As a result of extensive investigations with the purpose of settling the above-described problems, the inventors have found that a carnitine ester having high optical purity can be obtained in good yield by quaternarizing a  $\gamma$ -haloacetoacetic ester with trimethylamine followed by asymmetric hydrogenation in the presence of a ruthenium-optically active phosphine complex as a catalyst.

The present invention provides a process for preparing an optically active carnitine ester represented by formula (I):



wherein R represents a lower alkyl group; and X represents a halogen atom which comprises asymmetrically hydrogenating a  $\gamma$ -trimethylammonium-3-oxobutanoic ester halide represented by formula (II):



wherein R and X are as defined above,

in the presence of a ruthenium-optically active phosphine complex as a catalyst.


The compound represented by formula (II) which can be used in the present invention as a starting material can easily be synthesized by, for example, the process described in JP-A-51-63123.

In formula (II), the lower alkyl group as represented by R includes those containing from 1 to 4 carbon atoms, namely methyl, ethyl, butyl, n-propyl, isopropyl, isobutyl, and t-butyl groups; and the halogen atom as represented by X includes chlorine, bromine, and iodine atoms.

Examples of suitable ruthenium-optically active phosphine complexes which can be used for asymmetric hydrogenation include those represented by formulae (III), (V), (VI), and (VII):




wherein R<sup>1</sup>-BINAP represents a tertiary phosphine represented by formula (IV):



R<sup>1</sup> represents a hydrogen atom, a methyl group or a t-butyl group; Q represents a tertiary amine; when y is 0, then x represents 2, z represents 4, and p represents 1; and when y is 1, then x represents 1, z represents 1, and p represents 0,



wherein R<sup>1</sup>-BINAP is as defined above; Y represents ClO<sub>4</sub>, BF<sub>4</sub> or PF<sub>6</sub>; when u is 0, then v represents 1, and w represents 2; and when u is 1, then v represents 2, and w represents 1,



wherein R<sup>1</sup>-BINAP is as defined above; and R<sup>2</sup> represents a lower alkyl group (such as those containing from 1 to 4 carbon atoms) or a trifluoromethyl group, and



wherein R<sup>1</sup>-BINAP is as defined above; M represents Zn, Al, Ti or Sn; X' represents N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub> or CH<sub>3</sub>CO<sub>2</sub>; in the case that X' represents N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>, t is 2 and m is 1, and when M represents Zn, then k is 4, when M represents Al, then k is 5, and when M represents Ti or Sn, then k is 6; and in the case that X' represents CH<sub>3</sub>CO<sub>2</sub>, t is 1 and m is 2, and when M represents Zn, then k is 2, when M represents Al, then k is 3, and when M represents Ti or Sn, then k is 4.

The ruthenium-optically active phosphine complex represented by formula (III) can be obtained by the process disclosed in T. Ikariya et al., J. Chem. Soc., Chem. Commun., 922-924 (1985) and JP-A-61-63690.

The complexes represented by formulae (V) and (VI) can be obtained by the processes disclosed in JP-A-63-41487 and JP-A-62-265293, respectively.


The complex represented by formula (VII) can be obtained by starting with Ru<sub>2</sub>Cl<sub>4</sub>(R<sup>1</sup>-BINAP)<sub>2</sub>(NEt<sub>3</sub>)

(wherein Et represents, as it does hereinafter, an ethyl group), one of the complexes of formula (III), or  $\text{Ru}(\text{R}^1\text{-BINAP})(\text{OCOCH}_3)_2$ , one of the complexes of formula (VI). That is,  $\text{Ru}_2\text{Cl}_4(\text{R}^1\text{-BINAP})_2(\text{NEt}_3)$  or  $\text{Ru}(\text{R}^1\text{-BINAP})(\text{OCOCH}_3)_2$  is reacted with a Lewis acid selected from zinc chloride, aluminum chloride, titanium tetrachloride, and tin tetrachloride in a solvent, e.g., methylene chloride, at a temperature of from 10 to 25° C for a period of from 2 to 20 hours, and the solvent is removed from the reaction mixture by distillation, followed by drying to obtain the desired ruthenium-optically active phosphine complex as a solid.


Specific examples of the above-described ruthenium-optically active phosphine complexes which can be used in the present invention are shown below.













[wherein BINAP represents 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl, i.e., the tertiary phosphine of formula (IV) wherein  $\text{R}^1$  is a hydrogen atom]















[wherein T-BINAP represents 2,2'-bis(di-p-tolylphosphino)-1,1'-binaphthyl, i.e., the tertiary phosphine of formula (IV) wherein  $\text{R}^1$  is a methyl group]




[wherein t-Bu-BINAP represents 2,2'-bis(di-p-t-butylphenylphosphino)-1,1'-binaphthyl, i.e., the tertiary phosphine of formula (IV) wherein  $\text{R}^1$  is a t-butyl group]



(wherein t-Bu is a t-butyl group)



In carrying out the present invention, the compound of formula (II) is hydrogenated in an amphiprotic

solvent, e.g., methanol, anol, isopropanol, and acetic acid, at a temperature of from 15 to 100° C, preferably from 35 to 70° C, at a hydrogen pressure of from 10 to 150 kg/cm<sup>2</sup>, preferably from 25 to 50 kg/cm<sup>2</sup>, for a period of from about 2 to 40 hours, preferably from 5 to 30 hours. The ruthenium-optically active phosphine complex is used in an amount of from 1/50 to 1/5000 mole, preferably from 1/50 to 1/1000 mole, per mole of the compound of formula (II). The solvent is used in an amount of from 2 to 10 times, preferably from 3 to 5 times, the weight of the compound of formula (II).

In this reaction, the steric configuration of the compound of formula (I) can be controlled by selecting the BINAP moiety of the catalyst. For example, S-compounds can be obtained in using (R)-(+)-BINAP moieties, and R-compounds can be obtained in using (S)-(-)-BINAP moieties.

In particular, the S-compound of formula (I) can be lead to L-carnitine by hydrolysis.

After completion of the reaction, the reaction solution is concentrated under reduced pressure, and the residue is recrystallized to give the compound of formula (I). Alternatively, the residue is dissolved in water, the catalyst is removed by extraction from methylene chloride, etc., and the aqueous layer is concentrated to obtain a pure product. If desired, the catalyst may be reused for hydrogenation by removing the solvent from the extract of the catalyst and dissolving the residue in methanol or ethanol.

The present invention is now illustrated in greater detail with reference to Examples, but it should be understood that the present invention is not deemed to be limited thereto.

### EXAMPLE 1

In a 500 ml-volume stainless steel autoclave were charged 16.3 g (72.9 mmole) of ethyl  $\gamma$ -trimethylammonium-3-oxobutanoate chloride, 286 mg (0.169 mmole) of Ru<sub>2</sub>Cl<sub>4</sub>((-)-T-BINAP)<sub>2</sub>(NEt<sub>3</sub>), and 100 ml of ethanol, and a hydrogenation reaction was conducted at a hydrogen pressure of 50 kg/cm<sup>2</sup> and at 25° C for 18 hours under stirring. The ethanol was removed from the reaction mixture by distillation under reduced pressure of 20 mmHg to obtain 16.6 g of crude carnitine ethyl ester chloride.

$[\alpha]_D^{20}$ : -14.7° (c=1, H<sub>2</sub>O)

<sup>1</sup>H NMR  $\delta$  ppm: 2.28 (t, J=7.1Hz, 3H), 2.68 (q, 2H), 3.25 (s+m, 10H), 3.50 (d, 2H), 4.23 (dq, J=2.48Hz, 7.1Hz, 2H), 4.7 (m, 1H)

10 grams of the resulting crude carnitine ethyl ester chloride was dissolved in 20 ml of isopropyl alcohol at 50 to 52° C and then cooled to 5° C, followed by filtration to obtain pure carnitine ethyl ester chloride in a yield of 82%.

$[[\alpha]_D^{20}$ : -17.2° (c=1, H<sub>2</sub>O)].

The above-prepared crude carnitine ethyl ester chloride (10.6 g) was refluxed in 40 ml of water and 2 ml of concentrated hydrochloric acid for 2 hours, and the solution was concentrated to dryness at 60 to 65° C under reduced pressure of 10 mmHg to obtain 9.2 g of carnitine hydrochloride (optical yield: 85.4 %ee).

$[\alpha]_D^{25}$ : -20.23° (c=3.59, H<sub>2</sub>O)

<sup>1</sup>H NMR  $\delta$  ppm: 2.68 (q, 2H), 3.24 (s, 9H), 3.50 (d, 2H), 4.70 (m, 1H)

### EXAMPLE 2

In 500 ml-volume stainless steel autoclave were charged 16.3 g (72.9 mmole) of ethyl  $\gamma$ -trimethylammonium-3-oxobutanoate chloride, 328 mg (0.169 mmole) of Ru<sub>2</sub>Cl<sub>4</sub>((-)-BINAP)<sub>2</sub>(NEt<sub>3</sub>), and 100 ml of ethanol, and a hydrogenation reaction was conducted at a hydrogen pressure of 25 kg/cm<sup>2</sup> and at 50° C for 15 hours while stirring. The reaction mixture was distilled under reduced pressure of 15 mmHg to remove the ethanol to obtain 16 g of carnitine ethyl ester chloride.

$[\alpha]_D^{25}$ : -14.53° (c=1, H<sub>2</sub>O)

To the resulting carnitine ethyl ester chloride was added 50 ml of water, and subsequently 50 ml of methylene chloride was added thereto, followed by stirring. The oily layer was separated by liquid-liquid separation. To the aqueous layer was added 2 ml of concentrated hydrochloric acid, and the solution was heated at reflux for 2 hours, followed by concentration to dryness at 60 to 65° C under reduced pressure of 1 mmHg to obtain 9.0 g of carnitine hydrochloride (optical purity: 84 %ee).

$[\alpha]_D^{25}$ : -18.48° (c=4.00, H<sub>2</sub>O)

EXAMPLE 3

In 500 ml-volume stainless steel autoclave were charged 23.75 g (0.1 mole) of isopropyl  $\gamma$ -trimethylammonium-3-oxobutanoate chloride, 422 mg (0.25 mmole) of  $\text{Ru}_2\text{Cl}_4((-)\text{-BINAP})_2(\text{NEt}_3)$ , and 150 ml of isopropyl alcohol, and a hydrogenation reaction was conducted at a hydrogen pressure of 100 kg/cm<sup>2</sup> and at a temperature of 40 °C for 20 hours. The solvent was removed by distillation under reduced pressure of 10 mmHg to obtain 23.6 g of carnitine isopropyl ester chloride.

To the resulting carnitine isopropyl ester chloride was added 70 ml of water, followed by thoroughly stirring. After liquid-liquid separation, 3 ml of concentrated hydrochloric acid was added to the aqueous layer, and the solution was heated at reflux for 5 hours, followed by concentration to dryness at a liquid temperature of 60 to 65 °C under reduced pressure of 1 mmHg to obtain 16.7 g of carnitine hydrochloride (optical yield: 80 %ee).

$[\alpha]_D^{25}$  : -18.95° (c = 1, H<sub>2</sub>O)

EXAMPLE 4

In a 500 ml-volume stainless steel autoclave were charged 25.15 g (0.1 mole) of t-butyl  $\gamma$ -trimethylammonium-3-oxobutanoate chloride, 841 mg (1 mmole) of  $\text{Ru}((-)\text{-BINAP})(\text{OCOCH}_3)_2$ , and 100 ml of methanol, and a hydrogenation reaction was conducted at a hydrogen pressure of 70 kg/cm<sup>2</sup> and at a temperature of 25 °C for 40 hours. The solvent was removed by distillation under reduced pressure of 20 mmHg to obtain 25.5 g of carnitine t-butyl ester chloride. The product was treated in the same manner as in

Example 3 to obtain 17.7 g (90%) of L-carnitine hydrochloride (optical yield: 78.5 %ee).

$[\alpha]_D^{25}$  : -23.69° (c = 1.05, H<sub>2</sub>O)

EXAMPLE 5

In a 500 ml-volume stainless steel autoclave were charged 16.3 g (0.1 mole) of ethyl  $\gamma$ -trimethylammonium-3-oxobutanoate chloride, 355 mg (0.18 mmole) of  $[\text{Ru}((+)\text{-BINAP})\text{SnCl}_6]_2(\text{NEt}_3)$ , and 100 ml of ethanol, and a hydrogenation reaction was conducted at a hydrogen pressure of 100 kg/cm<sup>2</sup> and at 35 °C for 20 hours. The solvent was removed by distillation, and the residue was washed three times with 30 ml portions of methylene chloride, followed by drying at room temperature under reduced pressure of 5 mmHg for 17 hours to obtain 16 g of D-carnitine ethyl ester chloride (optical yield: 86 %ee).

$[\alpha]_D^{25}$  : +14.88° (c = 1.0, H<sub>2</sub>O)

EXAMPLE 6

The procedure of Example 5 was repeated, except for replacing  $[\text{Ru}((+)\text{-BINAP})\text{SnCl}_6]_2(\text{NEt}_3)$  with  $[\text{Ru}((-)\text{-BINAP})(\text{ClO}_4)_2]$  to obtain 16.1 g of L-carnitine ethyl ester chloride (optical yield: 83.8 %ee).

$[\alpha]_D^{25}$  : -14.5° (c = 1.0, H<sub>2</sub>O)

EXAMPLES 7 TO 15

Crude carnitine ethyl ester chloride was obtained in the same manner as in Example 1, except for altering the amount of the catalyst  $[\text{Ru}_2\text{Cl}_4((-)\text{-T-BINAP})_2(\text{NEt}_3)]$ , the kind of the solvent, the hydrogen pressure, the reaction temperature, and the reaction time as indicated in Table 1 below. The optical yields of the products are shown in the Table.


TABLE 1

Example No.	Catalyst/Substrate (mole/mole)	Solvent	Reaction Conditions			
			Hydrogen Pressure (kg/cm <sup>2</sup> )	Temp. (°C)	Time (hr)	Optical Yield (%ee)
7	1/230	ethanol	50	25	16	86.9
8	1/230	ethanol	50	50	4	84.6
9	1/230	ethanol	50	100	0.5	79.8
10	1/750	ethanol	50	80	3	78.0
11	1/750	ethanol	50	70	4	83.5
12	1/750	ethanol	50	60	5	85.0
13	1/750	ethanol	25	70	5	82.0
14	1/750	ethanol	10	70	6	82.0
15	1/230	acetic acid	50	50	6	89.8


As described above, the present invention provides a process for preparing an optically active carnitine ester which is an intermediate for obtaining pharmaceutically important carnitine. Using an easily available substrate and conducting asymmetrical hydrogenation in the presence of a ruthenium-optically active phosphine complex as a catalyst, the process of the present invention produces the desired compound in higher yield and at higher optical purity with high efficiency as compared with the conventional processes requiring complicated procedures and, thus, the process of the invention is of industrial advantage. Since a product of any desired isomerism can be arbitrarily obtained depending on the optical properties of the ruthenium-optically active phosphine complex used as a catalyst, it is expected to apply the present invention to a broadened range of use.

### Claims

1. A process for preparing an optically active carnitine ester represented by formula (I):




wherein R represents a lower alkyl group; and X represents a halogen atom, which comprises asymmetrically hydrogenating a  $\gamma$ -trimethylammonium-3-oxobutanoic ester halide represented by formula (II):




wherein R and X are as defined above,

in the presence of a ruthenium-optically active phosphine complex as a catalyst.

2. A process as claimed in Claim 1, wherein said ruthenium-optically active phosphine complex is selected from compounds represented by formula (III):




wherein R'-BINAP represents a tertiary phosphine represented by formula (IV):



(IV)


15  $R^1$  represents a hydrogen atom, a methyl group or a t-butyl group; Q represents a tertiary amine; when y is 0, then x represents 2, z represents 4, and p represents 0; and when y is 1, then x represents 1, z represents 1, and p represents 0;

compounds represented by formula (V):



20 wherein  $R^1-BINAP$  is as defined above; Y represents  $ClO_4$ ,  $BF_4$  or  $PF_6$ ; when u is 0, then v represents 1, and w represents 2; and when u is 1, then v represents 2, and w represents 1;

compounds represented by formula (VI):



25 wherein  $R^1-BINAP$  is as defined above; and  $R^2$  represents a lower alkyl group or a trifluoromethyl group; and compounds represented by formula (VII):



30 wherein  $R^1-BINAP$  is as defined above; M represents Zn, Al, or Ti or Sn;  $X^1$  represents  $N(C_2H_5)_3$  or  $CH_3CO_2$ ; in the case that  $X^1$  represents  $N(C_2H_5)_3$ , l is 2 and m is 1, and when M represents Zn, then k is 4, when M represents Al, then k is 5, and when M represents Ti or Sn, then k is 6; and in the case that  $X^1$  represents  $CH_3CO_2$ , l is 1 and m is 2, and when M represents Zn, then k is 2, when M represents Al, then k is 3, and when M represents Ti or Sn, then k is 4.

3. A process as claimed in Claim 1, 2 or 3, wherein said ruthenium-optically active phosphine complex is used in an amount of from 1/50 to 1/5000 mole per mole of the compound of formula (II).

35 4. A process as claimed in Claim 3, wherein said ruthenium-optically active phosphine complex is used in an amount of from 1/50 to 1/1000 mole per mole of the compound of formula (II).

5. A process as claimed in any preceding claim, wherein the hydrogenation is carried out at a temperature of from 15 to 100° C and at a hydrogen pressure of from 10 to 150 kg/cm<sup>2</sup> for a period of from about 2 to 40 hours.

40 6. A process as claimed in Claim 5, wherein the hydrogenation is carried out at a temperature of from 35 to 70° C and at a hydrogen pressure of from 25 to 50 kg/cm<sup>2</sup> for a period of from 5 to 30 hours.

7. A process as claimed in any preceding claim, wherein the hydrogenation is carried out in an amphiprotic solvent.

8. A process as claimed in Claim 7, wherein the solvent is methanol, ethanol, isopropanol, or acetic acid.

45 9. A process as claimed in Claim 7 or 8, wherein the solvent is used in an amount of from 2 to 10 times the weight of the compound of formula (II).

10. A process as claimed in Claim 9, wherein the solvent is used in an amount of from 3 to 5 times the weight of the compound of formula (II).



(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 89313429.6

(51) Int. Cl.<sup>5</sup>: **C07C 227/32, C07C 229/22, B01J 31/24**

(22) Date of filing: 21.12.89

(30) Priority: 22.12.88 JP 324651/88

(43) Date of publication of application:  
27.06.90 Bulletin 90/26

(84) Designated Contracting States:  
**CH DE FR GB IT LI NL**

(89) Date of deferred publication of the search report:  
22.11.90 Bulletin 90/47


(71) Applicant: **Takasago International Corporation**  
**19-22, Takanawa 3-chome Minato-ku**  
**Tokyo(JP)**

(72) Inventor: **Taketomi, Takanao**  
**112-7, Ookido-cho**  
**Chiba-shi Chiba(JP)**  
Inventor: **Sakaguchi, Toshiaki**  
**3-3-35, Minamihara**  
**Hiratsuka-shi Kanagawa(JP)**  
Inventor: **Kumobayashi, Hidenori**  
**1-4-39, Nakakaigan**  
**Chigasaki-shi Kanagawa(JP)**


(74) Representative: **Moore, Anthony John et al**  
**Gee & Co. Chancery House Chancery Lane**  
**London WC2A 1QU(GB)**

(54) **Process for preparing optically active carnitine ester.**

(57) A process for preparing an optically active carnitine ester represented by formula (I):



wherein R is lower alkyl and X is halogen, comprises asymmetrically hydrogenating a  $\gamma$ -trimethylammonium-3-oxobutanoic ester halide represented by formula (II):



wherein R and X are as defined above, in the presence of a ruthenium-optically active phosphine complex as a catalyst. An optically active carnitine ester of any desired isomerism can be obtained through simple operation in high yield at high optical purity. The substrate used is readily available.



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 89 31 3429

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,A	EP-A-0 339 764 (TAKASAGO) * Entire document * ---	1	C 07 C 227/32 C 07 C 229/22
A	TETRAHEDRON LETTERS, vol. 29, no. 13, 1988, pages 1555-1556, Pergamon Press plc, GB; M. KITAMURA et al.: "A practical asymmetric synthesis of carnitine" * Entire document * ---	1	
A	JOURNAL OF AMERICAN CHEMICAL SOCIETY, vol. 109, 1987, pages 5856-5858, American Chemical Society; R. NOYORI et al.: "Asymmetric hydrogenation of beta-keto carboxylic esters. A practical, purely chemical access to beta-hydroxy esters in high enantiomeric purity" * Entire document * ---	1	
A	EP-A-0 169 614 (CNRD) * Entire document * -----	1	TECHNICAL FIELDS SEARCHED (Int. Cl.5)  C 07 C 227/00 C 07 C 229/00
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 03-09-1990	Examiner WELLS A.G.
<b>CATEGORY OF CITED DOCUMENTS</b>  X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document  T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document			